

***Smeringopus butare* Huber, 2012**

Huber, B. A. 2012. Revision and cladistic analysis of the Afrotropical endemic genus *Smeringopus* Simon, 1890 (Araneae: Pholcidae). *Zootaxa* 3461: 1-138.

p. 93




562. *S. butare*, male dorsal view.

p. 94



FIGURES 572–583. *Smeringopus peregrinus* group, left male palps, prolateral and retrolateral views.
576–577. *S. butare*.

p. 96



FIGURES 592–607. *Smeringopus peregrinus* group, epigyna, ventral views and cleared female genitalia, dorsal views.
596–597. *S. butare*.




FIGURE 608. Known distribution of the *peregrinus* group. *S. butare* and *S. peregrinus* are represented by triangles in order to facilitate distinction from *S. peregrinoides*. Squares: further undescribed species.

Smeringopus butare new species


Figs. 562, 576–577, 596–597, 647–649, 663–666

Type. Male holotype from Rwanda, Butare [2°36'S, 29°44'E], vi.1971 (P. Nyalugaka), in MRAC (139117, part).

Etymology. The name is a noun in apposition, derived from the type locality.

Diagnosis. Distinguished from similar congeners by tip of procurus (distinctive membranous flap longer than sclerotized spine; Figs. 647, 648), and shape of embolus (very similar *S. peregrinus* but narrower in prolateral view

and without pointed process; Fig. 649); from *S. peregrinoides* also by longer and more downward directed cheliceral apophyses (similar *S. peregrinus*; cf. Figs. 614, 615) and absence of v-shaped or u-shaped structure frontally in female internal genitalia (Fig. 596; epigynum and cleared female genitalia otherwise similar *S. peregrinoides* and *S. kalomo*).



FIGURES 647–654. *Smeringopus butare* (647–649) and *S. katanga* (650–654). 647, 650. Left cymbia and procurus, retrolateral views. 648, 651–652. Left procurus, prolateral (648, 652) and dorsal (651) views. 649, 653. Left emboli, prolateral views. 654. Cleared female genitalia, dorsal view. Scale lines: 0.3 mm.

Male (holotype). Total body length 7.2, carapace width 2.2. Leg 1: 49.5 (13.1 + 0.9 + 13.1 + 19.7 + 2.7), tibia 2: 9.3, tibia 3: 6.8, tibia 4: 9.5; tibia 1 L/d: 55. Habitus as in Fig. 562. Carapace ochre-yellow with brown median and lateral marks, clypeus with pair of dark stripes, sternum brown with light marks near bases of coxae 2–4 and medially, legs light brown, femora and tibiae with lighter tips and subdistal dark rings, abdomen ochre-gray with distinct dorsal and ventral pattern. Distance PME-PME 185 μ m, diameter PME 170 μ m, distance PME-ALE 70 μ m, distance AME-AME 55 μ m, diameter AME 160 μ m. Ocular area slightly elevated, secondary eyes with ‘pseudo-lenses’; deep thoracic pit. Chelicerae with pair of distal apophyses as in *S. peregrinus* (cf. Figs. 614, 615; apophyses slightly shorter); each apophysis with one modified hair at tip (Fig. 663). Palps as in Figs. 576 and 577, coxa with distinct retrolateral apophysis and shallow wide furrow, trochanter barely modified, femur with retrolateral furrow with distinct rim proximally, procurus with distinctive distal elements (Figs. 647, 648), bulb with distinctively shaped prolateral process on embolus (similar *S. peregrinus* but narrower in prolateral view; Fig. 649). Legs without spines, few vertical hairs, with curved hairs ventrally on tibiae and metatarsi 1, retrolateral trichobothrium on tibia 1 at 3%; prolateral trichobothrium present on tibia 1. Gonopore with two epiandrous spigots (Fig. 664); ALS with eight spigots each (Fig. 665).

Variation. The color pattern is quite constant but variably distinct. Tibia 1 in 17 males: 9.3–18.1 (mean 13.8).


Female. In general similar to male; tibia 1 in 31 females: 10.0–15.7 (mean 11.9). Epigynum a simple plate mostly with poorly defined borders, with large semicircular pockets (Figs. 596, 666); internal genitalia as in Fig. 597 (very similar *S. peregrinoides* and *S. kalomo*).

Distribution. Known from Rwanda, Burundi, and eastern Congo D.R. (Sud-Kivu) (Fig. 608).

Material examined. RWANDA: Butare: type above; same data, 6♂4♀ in MRAC (139117, part); same data but vi.–vii.1971, 6♂3♀ in MRAC (140658 part), x.–xi.1970, 2♂15♀ + juvs (6 vials) in MRAC (137762 part, 137764 part, 137769, 137783, 137788, 137793); same locality, 23.xii.1979 (A. Vandenberghe), 1♀ 1 juv. in MRAC (153857). Astrida [=Butare, 2°36’S, 29°44’E], 30.vi.1952 (Laurent), 1♂1♀ 1 juv. in MRAC (73264–68). Near Lac Ihema [~1°55’S, 30°42’E], 14./18.vii.1969 (R. Kiss), 1♂ in MRAC (136301 part).

BURUNDI: *Bubanza Prov.*: “Crête Congo-Nil” [~3°S, 29.4°E], 2000 m a.s.l., iii.1967 (S. Ndani), 5♂6♀ (3 vials) in MRAC (132768–69, 132778). *Bururi Prov.*: Cabara [4°07’S, 29°31’E], Miombo woodland with *Brachystegia*, 850 m a.s.l., 18.iii.2002 (N. Benoit), 2♀ (2 vials) in MRAC (213925, 214163).

CONGO D.R.: *Sud-Kivu Prov.*: Bitale [2°11’S, 28°37’E], ~1700 m a.s.l., 30.–31.v.1949 (Laurent), 2♂ (2 vials) in MRAC (66508, 66512). Bukavu, Bitale, 19.iii.1950 (G. Marlier), 1♀ in MRAC (69051). Uvira [3°24’S, 29°08’E], vallée de lac Tanganika, 700 m a.s.l., “marais herbacé”, viii.1961 (R. Kiss), 1♂ in MRAC (120084); Uvira, 700 m a.s.l., vii.1961 (R. Kiss), 1♀ in MRAC (120048). Uvira, entre Kalundu et Kavimvira [~3°22’S, 29°09’E], vi.1961 (R. Kiss), 2♀ in MRAC (119923 part). Luhoho, Riv. Bunyakiri [2°05’S, 28°34’E], 1100 m a.s.l., 6.ix.1957 (E.S. Ross, R.E. Leech), 1♀ in CAS. Lwiro River, 47 km N Bukavu [~2°15’S, 28°50’E], 1950 m a.s.l., 15.xii.1957 (E.S. Ross, R.E. Leech), 1♀ 2 juvs in CAS. Irangi [1°54’S, 28°28’E], Luhoho River, 900 m a.s.l., 10.ix.1957 (E.S. Ross, R.E. Leech), 2♂1♀ in CAS. Irangi, vii.1964 (J. Bafort), 1♀ 1 juv. in MRAC (127421). Terr. Kalche, Maskele, 6 km from Irangi, “dans les creux des arbres vivantes en forêt primaire”, i.1962 (R. Kiss), 1♂1♀ in MRAC (121359).



FIGURES 663–672. *Smeringopus butare* (663–666) and *S. kalomo* (667–672). 663. Male cheliceral apophysis. 664. Male gonopore. 665. Male ALS. 666. Epigynum. 667. Male cheliceral apophysis. 668–669. Left procurus, retrolatero-dorsal and dorsal views. 670. Male gonopore. 671. Male ALS. 672. Epigynum. Scale lines: 10 μ m (663, 667), 20 μ m (671), 30 μ m (665), 60 μ m (668), 80 μ m (664, 669), 100 μ m (670), 200 μ m (666, 672).