


***Smeringopus hanglip* Huber, 2012**

Huber, B. A. 2012. Revision and cladistic analysis of the Afrotropical endemic genus *Smeringopus* Simon, 1890 (Araneae: Pholcidae). *Zootaxa* 3461: 1-138.

p. 45



FIGURES 243–255. *Smeringopus natalensis* group, habitus and male prosomata, oblique frontal views.

247–248. *S. hanglip*, male, dorsal and ventral views.


253. *S. hanglip*.

p. 47



FIGURES 268–277. *Smeringopus natalensis* group, left male palps, prolateral and retrolateral views.

270–271. *S. hanglip*.



FIGURES 278–297. *Smeringopus natalensis* group, epigyna, ventral views and cleared female genitalia, dorsal views.

290–291. *S.*

hanglip.




FIGURE 299. Known distribution of the *natalensis* group except *S. natalensis* (cf. Fig. 298). Squares: further undescribed species.


Smeringopus hanglip new species

Figs. 247–248, 253, 270–271, 290–291, 336–342, 350–356


Type. Male holotype from South Africa, Northern Province, Soutpansberg, 8 km NW Louis Trichard, Hanglip Forest, picnic area (~23°00'S, 29°53'E), 1440 m a.s.l., 30.xi.1996 (C.E. Griswold), in CAS.

Etymology. The name is a noun in apposition, derived from the type locality.

Diagnosis. Distinguished from most congeners (except *S. lydenberg*) by two black lines ventrally on abdomen (versus three; Fig. 248); from similar congeners (*S. lydenberg*, *S. ndumo*, *S. mlilwane*) by shapes of bulbal processes (Figs. 338, 339); from other close relatives by low process near palpal tarsal organ (Fig. 336), ventrally strongly curved procursus (Figs. 271, 336), and prolateral process on procursus tip (Fig. 337).



FIGURES 336–342. *Smeringopus hanglip*. 336. Left cymbium and procursus, retrolateral view. 337. Left procursus, dorsal view. 338–339. Left bulbal processes, prolateral and dorsal views. 340–341. Male chelicerae, frontal and lateral views. 342. Cleared female genitalia, dorsal view (arrow points to internal pocket). Scale lines: 0.3 mm (336–339), 0.5 mm (340–342).



FIGURES 343–356. *S. hanglip* (350–356). 343. Left procursus tip, retrolateral view. 344. Left palpal femur, ventral view. 345. Male cheliceral apophysis. 346. Tarsal pseudosegments. 347. Male gonopore. 348. Cleared female genitalia, dorsal view. 349. Detail of pore plate. 350. Left procursus, dorsal view. 351. Male cheliceral apophysis. 352. Bulbal processes. 353. Male ALS and PMS. 354. Male gonopore. 355. Epigynum. 356. Female ALS. Scale lines: 20 μm (345, 351), 30 μm (346, 356), 40 μm (353), 50 μm (349), 60 μm (343), 80 μm (347), 100 μm (344, 352), 200 μm (348, 350, 354), 300 μm (355).

Male (holotype). Total body length 9.5, carapace width 3.3. Leg 1: 77.8 (20.4 + 1.5 + 19.6 + 33.2 + 3.1), tibia 2: 14.5, tibia 3: 11.3, tibia 4: 14.8; tibia 1 L/d: 59. Habitus as in Figs. 247 and 248. Carapace ochre-yellow with

distinct dark pattern (median and lateral bands, no submarginal marks), clypeus with pair of indistinct dark stripes, sternum posterior half brown, legs with indistinct darker rings subdistally on femora and tibiae, abdomen dorsally with distinct dark pattern, ventrally with two dark lines. Distance PME-PME 240 μm , diameter PME 240 μm , distance PME-ALE 90 μm , distance AME-AME 70 μm , diameter AME 205 μm . Ocular area slightly elevated, secondary eyes with very indistinct 'pseudo-lenses'; deep thoracic pit. Chelicerae as in Figs. 340 and 341; with pair of small distal apophyses; each apophysis with modified hair at tip (Fig. 351). Palps as in Figs. 270 and 271, coxa without retrolateral apophysis, trochanter barely modified, femur with deep and wide retrolateral furrow with distinct rim proximally, cymbium with indistinct projection near tarsal organ (Fig. 336), procurus ventrally strongly curved (Figs. 271, 336), with prolateral process at tip (Figs. 337, 350), bulb with three distinctively shaped processes (Figs. 338, 339, 352). Legs without spines, few vertical hairs, with curved hairs ventrally on tibiae and metatarsi 1 and 2, retrolateral trichobothrium on tibia 1 at 1.5%; prolateral trichobothrium present on tibia 1. Gonopore with two epiandrous spigots (Fig. 354); ALS with eight spigots each (Fig. 353).

Variation. Tibia 1 in 6 other males: 14.0–19.2 (mean 16.5). In southern specimens (Magoebaskloof and George's Valley) the two dorsal processes of the bulb are slightly closer together.

Female. In general similar to male; tibia 1 in 17 females: 12.1–18.8 (mean 15.7). Epigynum a simple plate without pockets (Figs. 290, 355), laterally whitish, not clearly distinguishable from close relatives (*S. lydenberg*, *S. mlibwane*); internal genitalia as in Figs. 291 and 342 (also similar to close relatives, with internal pockets). ALS as in male (Fig. 356).

Distribution. Known from several localities in northeastern South Africa (Fig. 299).

Material examined. SOUTH AFRICA: *Limpopo*: Hanglip Forest: 1♂ holotype above; same data, 2♂7♀ in CAS. Soutpansberg, Entabeni Forest, ~20 km N Levubu (22°59'S, 30°17'E), 1360 m a.s.l., 1.–2.xii.1996 (C.E. Griswold), 3♂5♀ in CAS. Magoebaskloof Hotel, 30 km SSW Tzaneen (~23°53'S, 30°00'E), 22.–23.xi.1996 (C.E. Griswold), 3♀ 1 juv. in CAS. 28 km SSW Tzaneen, 8.6 km from Magoebaskloof Hotel, forest on Magoebaskloof trail (~23°50'S, 29°59'E), 1800 m a.s.l., 22.–23.xi.1996 (C.E. Griswold), 1♀ 4 juvs in CAS. Magoebaskloof, Hideaway Farm, forest, pump house, near stream, 11.viii.1997 (R. Jocqué), 1♂1♀ in MRAC (206534); same data but 1700 m a.s.l., night catch, 1♀ 2 juvs in MRAC (206529). George's Valley [~23°57'S, 30°01'E], 23.iii.2001 (G. Binford), 1♂2♀ in ZFMK (Ar 8508).